



Supporting informed-decision making in efforts to improve screening participation: implications for 'hard to reach' recruitment strategies

Dr Christine Campbell, University of Edinburgh

World Cancer Congress, Montreal
Thursday 30th August 2012



The Scottish School of Primary Care



Overview of this presentation

1. The challenge of inequitable participation in cancer screening programmes
2. Informed choice & use of decision aids in cancer screening
3. Public preferences regarding level of recommendation – result from a UK survey
4. Concluding thoughts

(my focus will be on bowel cancer screening, and wrt population screening: the issues are just as pertinent for cervical and breast screening, and other screening contexts)

UK Bowel Screening Pilots

Socio-economic gradient
1 least deprived
5 most deprived



Minority ethnic populations

Barriers to participation in bowel screening

Generic

- Lack of clinical support (esp primary care provider)
- Fear (of cancer, treatment, colostomy)
- Lack of understanding of nature of screening
- Shame (embarrassment)
- Perception of personal risk
- Screening in absence of symptoms
- Conflicting priorities

Cultural

- Taboo around faecal matter
- Fatalism

Process

- Understanding of process
- Storage & Hygiene
- Use of medical terms
- Health literacy

Health System

- Cost – medical coverage
- Access

Strategies to address low participation

- **Screening modalities** – e.g. one-sample FIT versus three-sample guaiac FOBT; HPV self-sample for cervical screening
- **Improved access** - screening facility hours, location, costs
- **Adapting recruitment materials** -targeted & tailored materials for specific populations
- **‘pre-notification’ letters** (improved uptake by >5%) and **reminders** (improved uptake by 8%)
- **Primary care engagement** - endorsement of invitations; ‘Local Champion’ role; more extensive feedback on participation status; practice-based promotion of screening

Informed choice in cancer screening

- In order to **maximise the public health benefits** of screening **uptake must be high**
- Current important emphasis on patient involvement in health care; **informed choice principles**
- The UK's Department of Health Improving outcomes: a strategy for cancer – *'To empower the greatest number possible from all groups and communities... to make an informed choice to participate in cancer screening'*
- **Potential tension** between personal autonomy and public health benefit.

Rapid review: informed choice & cancer screening

- Five systematic reviews of informed choice and decision-making
 - Although most people value information on the limits of screening, provision of information and education alone are not sufficient to facilitate an informed choice
 - An individual's personal experience, values and health and social contexts are equally or more important
 - Few interventions are sufficiently grounded in theoretical models or conceptual frameworks

What about decision aids for cancer screening?

- Seven systematic reviews of decision aids
 - Good evidence that **DAs incorporating knowledge and patient values are effective in increasing informed decision-making**, reducing decisional conflict, and reducing anxiety
 - DAs do result in increased informed choice and decreased PSA testing; **data inconclusive for screening for other cancers**
 - Some evidence DAs can increase patient/physician communication (USA)


Rapid review –recent RCTs of DAs

Reference	Country	Details	Key findings
Smith et al BMJ 2010	Australia	Colorectal cancer screening for adults with low educational attainment ; extensive decision aid booklet	Increased knowledge, less positive attitude, increased proportion making an informed choice; significantly lower participation in bowel cancer screening in intervention arm
Steckelberg et al BMJ 2011	Germany	Colorectal cancer screening, brochure with comprehensive evidence-based risk information	Increased the proportion who indicated an informed choice; no difference in combined actual and planned uptake
Miller et al Am J Prev Med 2011	USA	Colorectal cancer screening - Web-based multi-media patient decision aid, designed for mixed-literacy abilities, in low SES population	DA significantly increased readiness to receive screening; non- sig increase in tests ordered and completed; results similar across literacy levels

- These examples of decision aids in colorectal cancer screening show that facilitating increase
- These examples of decision aids in colorectal cancer screening show that facilitating increase

“If interventions that foster informed decision-making reduce uptake of bowel cancer screening, then interventions that prioritise the soundness of individual patient decisions may work at cross-purposes with the overarching programme goal of reducing population cancer burden”

A fine balance



Promoting equitable uptake


Promoting informed decision-making

What do the public want?

- A recent UK study examined current attitudes towards and understanding of cancer screening
- Allowed us to explore public preferences for levels of recommendation about bowel cancer screening
- Carried out in 2011, random location sampling across the UK with quotas for gender, working status

Aims and objectives

- To assess public preferences for **the level of recommendation** provided by the National Health Service for the bowel screening programme in the UK
- To measure the public's desire about **the quantity of information** about the risks and benefits of screening
- To explore differences by socio-demographic characteristics



84% England; 11% Scotland; 5% Wales

4% minority ethnic origin


50 to 80 years (*Mean age*: 64.2)

AB: 20%, C1: 23%, C2: 20%, DE: 37%

62% married, 29% wid/div/sep, 9% single


52% female and 48% male

Recommendation vs. no recommendation from the NHS



- Over 80% preferred a clear recommendation to participate from the NHS
- Only 15% wanted information with no recommendation
- **No association between SES and recommendation preference**
- Men more likely than women to want a strong recommendation

Desire for benefit and risk information



Most want to receive **all** information available: even if they report not reading previous leaflets in full

Participants with lower SES had **lower understanding of the purpose** of screening, **wanted less information on benefits and harms** of screening;

Concluding thoughts...

- Imperative that researchers and practitioners rigorously apply informed choice principles when designing and implementing interventions for 'hard to reach' groups
- Autonomous decision-making does not preclude provider input: the survey suggests that the public would welcome a recommendation –not as an alternative to information, but as an *adjunct* to it
- Consistent with 'consider an offer' approach(Entwistle ,BMJ 2009)
- Those with lower SES wanted a recommendation to the same extent as higher SES, but placed less importance on receiving full information on benefits and harms: consistent with previous findings that barriers are often elsewhere
- Role for primary care providers in supporting invitees through a decision-making process: factors such as co-morbidities can be discussed before a negotiated 'recommendation' can be reached

Acknowledgements

- University of Edinburgh: Prof David Weller, Vicky Hammersley, Roma Robertson, Lucine Techer, Fay Crawford, Allison Worth
- Bowel Screening Programmes: Prof Bob Steele, Prof Julietta Patnick , Dr Steve Smith
- University College London: Jo Waller, Alice Simon, Christian von Wagner, Craig Jones, Ana Macedo, Jane Wardle
- Funders: Cancer Research UK, National Cancer Institute (USA), Scottish School of Primary Care



Ca-PRI Network



- An open, multidisciplinary network for researchers in primary care and cancer
- Promotes greater international collaboration in this field by supporting networking and arranging a yearly conference
- Focus is on the role of primary care throughout the cancer journey, from prevention to palliative care
- Website: <http://www.ca-pri.com/>

Ca-PRI 6th Annual Conference

will be held 15th - 16th April 2013

University of Cambridge, UK

Reserve the dates

(The conference will overlap with the UK's National Awareness and Early Diagnosis Initiative (NEADI) on the 16th/17th April)